



## CERTIFICATE OF ANALYSIS

Product BIOTINYLATED ANTI-CHICKEN IgG (H+L)  
\*\*AFFINITY PURIFIED\*\*

Catalog BA-9010

Amount 1.5 mg biotinylated IgG

Concentration 1.5 mg/ml

Lot No. X0809

Produced in goat

Solution 10 mM phosphate, pH 7.8, 0.15 M NaCl, 0.08% sodium azide. Immunohistochemical grade bovine serum albumin has been added for stabilization.

Storage Refrigerate; for long-term storage, aliquots may be stored frozen.

Immunoglobulin biotinylated 99%

Antibody Activity after biotinylation (by solid phase IgG binding assay) 97%

Solid-phase enzyme immunoassay of cross-reactivity:

<u>Protein</u>	<u>Percent Cross-reactivity</u>
rabbit IgG	3%
dog IgM, sheep IgG	1%
bovine IgG, bovine IgM, cat IgG, dog IgG, goat IgG, guinea pig IgG, hamster IgG, horse IgG, human IgG, human IgM, mouse IgG, rabbit IgM, rat IgG, rat IgM, Rhesus monkey IgG, sheep IgM, swine IgG, swine IgM	<1%

If this biotinylated antibody is to be used in tissues, which may contain cross-reacting endogenous immunoglobulins, dilution of this biotinylated antibody may be made in buffers containing 2% normal serum from the same species as the tissue.

The recommended concentration range for use is 2-10 µg/ml.